Repositioning the Hemodynamically Unstable "Do Not Turn" Patient

- Even if patient cannot tolerate a full turn, most patients can still be turned to a lower angle (micro-turn)
- Reposition the head and limbs, consider passive ROM hourly
- Almost all patients can still have their heels floated. If this is impossible, apply Mepilex heel dressing (#189956)until able to use offloading boots or pillow


- Please note: A Bedrest order is not the same thing as a Do Not Turn order
- Be aware of all lines and devices attached to patient, ensure none are under the patient (swipe for debris)
- Use a Mepilex dressing or fluidized positioner pillow (Z-Flow), to cushion or offload
- When attempting to resume turning, begin slowly, small serial turns

Orders:


Restrictions: Do not turn

Order sentences for: Restrictions

(None)
T;N, Do not allow patient to turn to operative side
T;N, Do not shave chest hair
T;N, Keep head of bed flat
T;N, No smoking privileges
T;N, Restriction Type: No Intramuscular Injections



Activity: Bedrest



This will fire a consult to the WOCN to make aware of pt's increased risk

_	Consult to Infection Prevention	Ordered	08/03/20 9:42:50 Consult placed due to patient meeting criteria for PUI	SYSTEM
	Consult to Wound Ostomy Continence Team (Inpatient)	Ordered	09/01/20 14:07:21 Consult placed by rule due to the Restriction order: Do Not Turn being initiated on the patient.	SYSTEM
\checkmark	Home Care Assessment - Case	Ordered	08/03/20 8:41:13, "BOOST: Prior Hospitalizations has occurred due to 2 or more inpatient/observation hospitalizations or 2 or m	SYSTEM